



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$$\Sigma my^2 = \rho \sin^3 \beta \int_{-\frac{1}{2}c}^{\frac{1}{2}c} \int_0^c y^2 dx dy = \frac{1}{8} mc^2 \sin^2 \beta.$$

$$\therefore \tan 2\theta = \frac{4 \sin 2\beta}{1 + 4 \cos 2\beta}.$$

$$\text{But } 2\theta = \beta + 2\varphi. \quad \therefore \tan 2\varphi = \frac{1}{3} \tan \beta.$$

$$A \cos^2 \theta + \beta \sin^2 \theta = \frac{1}{2} m (c^2 + 4c^2 \cos^2 \beta), \quad A \sin^2 \theta + \beta \cos^2 \theta = \frac{1}{2} mc^2 \sin^2 \beta.$$

$$\therefore A = \frac{1}{24} mc^2 [5 + \sqrt{(17 + 8 \cos 2\beta)}], \quad B = \frac{1}{24} mc^2 [5 - \sqrt{(17 + 8 \cos 2\beta)}].$$

$$c^2 = a^2 + b^2, \quad \cos \beta = \frac{a^2 - b^2}{a^2 + b^2}, \quad \cos 2\beta = \frac{(a^2 + b^2)^2 - 8a^2 b^2}{(a^2 + b^2)^2}.$$

$$\therefore A = \frac{1}{24} m \{ 5a^2 + 5b^2 + \sqrt{[25(a^2 + b^2) - 64a^2 b^2]} \}.$$

$$B = \frac{1}{24} m \{ 5a^2 + b^2 - \sqrt{[25(a^2 + b^2) - 64a^2 b^2]} \}.$$

106. Proposed by J. E. CRAIG, A. B., New Germantown, N. J.

The centers of the two wheels of a bicycle are three feet apart.

(1) If a rider wishes the rear wheel to trace a circle 14 feet in diameter, what must be the diameter of the circle traced by the front wheel?

(2) If the rider weighs 120 pounds, and his center of gravity is 3 feet from the ground, at what angle must he lean to make one revolution of the circle every 3 seconds?

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

(1) The line that is tangent to the hind wheel's path passes through the center of the front wheel, caused by the rigid frame work.

$$\therefore R = \sqrt{(9 + 49)} = \sqrt{58} = 7.615773 \text{ feet.}$$

$$2R = \text{diameter traced by front wheel} = 15.231546 \text{ feet.}$$

(2) Since the man must be in equilibrium we do not use his weight, and we may regard him as moving around in the plane of the track. Suppose he is midway between the wheels.

Let  $\theta$  be his inclination to the vertical,  $g$  = gravity,  $f = v^2/r$  = centrifugal force. Then  $g \sin \theta$  = component of gravity perpendicular to man's direction,  $f \cos \theta$  = component of centrifugal force perpendicular to same direction. When in equilibrium these forces equalize each other.

$$\therefore g \sin \theta = f \cos \theta = (v^2/r) \cos \theta.$$

$$\therefore \tan \theta = v^2/gr.$$


$$r = \sqrt{(49 + \frac{9}{4})} = 7.1589 \text{ feet.} \quad v = 14.3178\pi/3 \text{ feet} = 14.9936 \text{ feet per second.}$$

$$\therefore \tan \theta = \frac{(14.9936)^2}{32\frac{1}{2} \times 7.1589} = .97625.$$

$$\therefore \theta = 44^\circ 18' 41''.$$


107. Proposed by M. E. GRABER, Student, Heidelberg University, Tiffin, O.

Two particles attracting each other inversely as the square of their distances apart, are constrained to move in straight lines which intersect each other at right angles. How long will it take for the particles to meet and how far does each particle move?



Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Let  $OA=a$ ,  $OB=b$ ,  $OD=u$ ,  $OC=x$ ,  $m$ =mass of particle at  $C$ ,  $n$ =mass of particle at  $D$ ;  $A$ ,  $B$  the positions of the particles at the beginning of motion;  $C$ ,  $D$  their positions at any time  $t$ ;  $v$ ,  $v_1$  their velocities at  $C$ ,  $D$ . Then the equations of motion are



$$\text{for } C, \frac{d^2x}{dt^2} = \frac{nx}{(x^2+u^2)^{\frac{3}{2}}} = F, \text{ since } s=x,$$

$$\text{for } D, \frac{d^2u}{dt^2} = \frac{mu}{(x^2+u^2)^{\frac{3}{2}}} = f, \text{ since } \sigma=u. \text{ But } v=Ft, v_1=ft.$$

$$\therefore v_1 = \frac{mu v}{nx}. \text{ Also } \frac{a-x}{v} = \frac{b-u}{v_1} = \frac{(b-u)nx}{mu v}.$$

$$\therefore u = \frac{bnx}{am-mx+nx} \dots (1). \quad \therefore \frac{d^2x}{dt^2} = \frac{n(am-mx+nx)^3}{x^2[a^2n^2+(am-mx+nx)^2]^{\frac{3}{2}}} \dots (2).$$

From (1),  $u=0$  when  $x=0$ . Therefore, the particles both arrive at  $O$  at the same time. Hence  $C$  moves over distance  $a$ , and  $D$  moves over distance  $b$  before they meet. The time is found by integrating (if possible) (2) twice.

$$\text{If } m=n, \frac{d^2x}{dt^2} = \frac{a^3m}{x^2 \sqrt{[(a^2+b^2)^3]}}.$$

$$\therefore \left(\frac{dx}{dt}\right)^2 = \frac{2a^3m}{(a^2+b^2)^{\frac{3}{2}}} \left(\frac{1}{x} - \frac{1}{a}\right) = \frac{2a^2m}{(a^2+b^2)^{\frac{3}{2}}} \cdot \frac{a-x}{x}.$$

$$\therefore t = \frac{(a^2+b^2)^{\frac{3}{2}}}{a\sqrt{2m}} \int_0^a \sqrt{\frac{x}{a-x}} dx = \frac{\pi(a^2+b^2)^{\frac{3}{2}}}{2\sqrt{(2m)}}.$$

$$\text{If } a=b \text{ and } m=\text{unity}, t=\pi(\frac{1}{2}a^2)^{\frac{3}{2}}.$$

#### AVERAGE AND PROBABILITY.

92. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in Irving College, Mechanicsburg, Pa.

A circular field, radius  $r$ , is divided into four *equal* parts, by concentric circles and three concentric rings. From the center of this field are fired *at random*, and with such a velocity as not to produce a range greater than the radius of the field,  $m=1000$  projectiles of the *same* kind. How many projectiles should have fallen into each one of these four equal parts of the field?

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

The range  $= (v^2/g)\sin 2\theta$ , where  $\theta$  = angle of elevation. Greatest range  $= v^2/g = r$ .